

**SUSTAINABLE DEVELOPMENT OF SOUTHEAST ASIA'S MARINE
ECOSYSTEMS – CLIMATE CHANGE CHALLENGES
AND MANAGEMENT APPROACHES**

Loke Ming Chou

Department of Biological Sciences,
National University of Singapore,
Blk. S3, 14 Science Drive 4, Singapore 117543.
Email: db sclm@nus.edu.sg, Tel: (65) 6516 2696, Fax: (65) 6779 2486

ABSTRACT. *Southeast Asia is recognized as a global marine biodiversity hotspot because of its rich marine natural heritage. The valuable benefits of its marine ecosystem services however, have been eroding since the mid-1960s as heavy anthropogenic pressure driven by rapid economic growth and a fast-expanding population resulted in extensive ecosystem degradation and environmental quality decline. Awareness of the erosion of coastal and marine ecosystem integrity and appreciation of ecosystem services loss in terms of monetary value increased over the past few decades but management response has remained reactive rather than proactive. As human activity causes ecosystem resiliency loss at local scales, climate change impacts operating at the global scale can exacerbate biodiversity loss at further expense to the already compromised ecosystem resiliency. As warming augments, equatorial species already at high temperature tolerance limits will begin migrating to higher latitudes. Ecosystem integrity will be further reduced as species-specific range shifts occur at varying temporal and spatial scales and rates. Trophic disruptions, breakdown of species interactions and interference of ecosystem processes are likely to result in major modification of present community structures with overall biodiversity simplification challenging ecosystem integrity even further. Other relevant climate change impacts such as increased frequency of extreme weather, sea-level rise, and ocean acidification will also affect marine biodiversity. Such impacts can be seen from recent events that depict climate change impact scenarios. With many uncertainties, what may emerge as the most relevant mitigation priority is to reduce present anthropogenic pressures that will allow ecosystem integrity to improve and maintain some resiliency to the impacts of climate change. There are choices between ecosystem management, degradation and restoration. Case studies indicate possible significant and sustained economic returns from adequately managed marine ecosystems compared to irreversible loss from damaged ones. Habitat restoration initiatives have increased in the past two decades in response to widespread degradation although they are costlier and face limitations. Protection and management are the preferred options if ecosystem service benefits are to be optimized. Healthy, resilient ecosystems are also the best investment against the emerging threats of climate change and natural disasters.*

KEYWORDS. Marine biodiversity, climate change, impacts, Southeast Asia.

INTRODUCTION

Southeast Asia stretches over the equatorial belt between latitudes 21⁰N to 12⁰S and longitudes 93⁰E to 141⁰E. The surface area of all its seas combined total 9 million km² representing just 2.5% of earth's ocean surface but supports almost 30% of the world's coral reefs (Burke *et al.*, 2011), one third of the world's mangroves (FAO, 2007) and 17% of the

world's seagrass area (Green & Short, 2003). The region harbours a great variety of coastal and marine ecosystems, known throughout the world for its high species richness (Chou, 1996). The island-filled region is located in between the Asian and Australian continents and between the Pacific and Indian Oceans. The bathymetry is a mix of shallow continental shelves, continental slopes, deep basins, trenches and troughs.

Indonesia and the Philippines (the world's two largest archipelagos) have over 25,000 islands and almost all the region's countries along the Asian continent possess extensive coastlines and numerous offshore islands, most of which are coral or volcanic. The combined coastline length of 92,451 km amounts to approximately 16% of the world's total and this extensive coastline supports a wide variety of coastal and marine ecosystems. Coastal features such as cliffs, coves, beaches (rocky, sandy, muddy), deltas, spits, dunes and lagoons provide the potential of harbouring high species richness. The high diversity marine habitats are favoured by the tropical climate and heavy precipitation that transports nutrients from land to sea.

The high nutrient content of Southeast Asian seas is facilitated by the scattering of numerous islands, each contributing terrestrial inputs to the marine system. The warm tropical climate further contributes to boosting primary productivity, enabling mangrove forests to reach their maximum development and greatest luxuriance in parts of the region (Rao, 1986). A comparative review of mangrove ecosystem productivity based on different parameters such as phytoplankton production, primary production, benthic primary production and total litter production showed that the region's mangrove forests maintained the highest values than elsewhere in the world (Singh *et al.*, 1994)

Rapid industrialisation and strong economic development combined with the fast expanding human population exerted high pressure on marine biodiversity through habitat loss and degradation, pollution and overexploitation (UNEP/COBSEA, 2010). Economic growth generated mainly by industrialisation and international trade remained high during the 1980s and early 1990s (JEC, 2000). Evidently, economic development remained the clear focus of Southeast Asian states with the pace intensifying from the 1960s (UNEP/COBSEA, 2010). The link between biodiversity and climate change has been clearly established (IPCC, 2007). While climate change will affect biodiversity, the latter has a role in mitigating climate change influences (Djoghlaif & Ganapin Jr., 2010). The region's marine biodiversity is already under high threat from anthropogenic impacts that compromise its ability to provide ecosystem services that can at the same time mitigate climate change impacts.

ECOSYSTEM RICHNESS

The region's geomorphologic and oceanographic features together with the tropical weather make the seas highly productive and supportive of rich and extensive marine habitats. Recognised as the faunistic centre of the entire Indo-Pacific (IUCN/UNEP, 1985), species richness of corals and other reef-associated flora and fauna is the highest throughout the world making the region a global hotspot for coral reefs (Kelleher *et al.*, 1995). It has about 80% of the world's hard coral species (Spalding *et al.*, 2001), 60% of mangrove species (FAO, 2007) and 50% of seagrass species (Green & Short, 2003).

A major significance is that many taxa are present only in the region and their loss translates to global extinction. This species uniqueness has to be guarded against further loss. Investigations into population genetics of reef organisms such as corals (Knittweis *et al.*, 2009), fish (Lourie & Vincent, 2004; Timm & Kochzius, 2008), crustaceans (Barber *et al.*, 2006), molluscs (Kochzius & Nuryanto, 2008) and echinoderms (Kochzius *et al.*, 2009) indicate high levels of genetic structuring with distinct signatures from the Indo-Pacific.

A high proportion of coral, fish, gastropod and lobster species have restricted geographic ranges (Roberts *et al.*, 2002) and remain at high risk of extinction from localized reef degradation. Bellwood and Meyer (2009) demonstrated that the recognized (but arbitrarily defined) hotspots of the Indo-Australian Archipelago do not necessarily support high numbers of endemics or serve as a speciation source, and suggested that conservation of areas beyond these hotspots is also important. Weeks *et al.* (2010) also recognised from their assessment of community-based Marine Protected Areas (MPAs) in the Philippines that these small protected areas should be supplemented by larger no-take reserves for biodiversity conservation targets to be more effectively met.

HUMAN IMPACTS ON ECOSYSTEM SERVICES

Marine biodiversity of the region has suffered significant loss and degradation (UNEP/COBSEA, 2010). Anthropogenic impacts are significant and increasingly depressing delivery of ecosystem services that are essential to the well-being of human society and national economies (UNEP, 2001; PEMSEA, 2003; MEA, 2005; ASEAN, 2006; UNEP/COBSEA, 2010). Human pressure on coastal and marine ecosystems is immense considering that 85% of the region's population is located within 100km of the coast, when the world's average coastal population is 40% (UNEP/COBSEA, 2010). It is estimated that 95% of Southeast Asia's coral reefs are at risk from local threats (Burke *et al.*, 2011). Long-term decline of Indo-Pacific reefs is evident with the proportion of reefs having at least 50% live coral cover falling from 66% in 1980 to 4% in 2004 (SCBD, 2010). Signs of decline in the rate of mangrove loss have emerged for the rest of the world but not in Asia, which has the greater proportion of the ecosystem (SCDB, 2010).

Sustained economic growth over the past four decades has helped countries in the region to improve national Gross Domestic Product and reduce poverty (PEMSEA, 2003). Most of the economic activity has been focused on coastal areas and a large proportion of the region's population is dependent on the coastal and marine environment and their resources (UNEP/COBSEA, 2010). This growth comes at the expense of ecosystem services that are lost from degraded habitats. The estimated annual value of ecosystem services from the region's coral reefs is US\$112.5 billion (PEMSEA, 2003). The annual total economic value per hectare of habitats in the South China Sea was estimated at US\$2,872 for mangroves, \$1,543 for coral reefs, \$1,182 for seagrass and \$295 for wetlands (UNEP, 2007).

CLIMATE CHANGE IMPACTS

There is increasing evidence that climate change can cause significant impacts to biodiversity (IPCC, 2007). Climate change exposes marine biodiversity to a wide range of impacts that include sea level rise, elevated sea temperature, and increased frequency of extreme weather and intensification of water column stratification (Chou, 1994). These impacts generate accompanying environmental effects on coastal and marine systems through coastal erosion, sudden salinity fluctuation, increased sedimentation, nutrient loading, salt water intrusion, coastal inundation, and changes in coastal geomorphology and circulation patterns (Chou, 1992). While it is not easy to isolate climate change impacts on natural systems from adaptation and other drivers not directly related to climate change, it is possible to get some idea of climate change impacts from existing scenarios that simulate climate change conditions (Chou, 2010).

Impacts on biodiversity can occur at the species and community levels. At the species level, physiological constraints limit the individual's tolerance to environmental changes. At the community level, changes in the community structure alter the ecosystem functioning of the habitats. A review of the implications of expected climate change impacts on natural coastal ecosystems in the region has been made by Yap (1994). Climate change impacts on species will vary. Some species can move to new areas with more suitable conditions, while sessile and sedentary ones will have to cope or perish. Many are not expected to adapt to the rate and intensity of projected climate change scenarios and risk extinction (SCBD, 2010). In the most simplistic sense, a shift to the higher latitudes is expected of fish and other pelagic species resulting in a possible decrease in equatorial biodiversity. However, large changes in community structure will take place as intact communities fragment from the departure of some species resulting in trophic cascade modification and ecological imbalance. The same effect will be seen at sites receiving immigrating species.

Rising sea levels inundating low-lying coastal plains can overwhelm adaptation response of coastal biomes and temperature elevation is expected to elicit physiological and behavioural responses from species that could be detrimental to entire biological communities and ecosystem integrity. Further warming of estuarine and near-shore habitats may make them inhospitable to species that already live close to the upper temperature tolerance limit. Increased precipitation will also test the tolerance limits of these species.

Ocean acidification from increased dissolved carbon dioxide is detrimental to many marine species such as corals, shellfish and plankton. Coral reefs in particular are highly vulnerable to lowered ocean pH and the region, with its high proportion of the world's reefs will suffer a greater loss. Should atmospheric CO₂ concentrations reach 500 ppm after 2050 as projected by Rogeli *et al.* (2009), coral growth will be arrested as dissolution takes place (Silverman *et al.*, 2009). The loss of coral reefs will be significant to the region where millions of people depend on it for subsistence. Reefs subjected long-term to lowered pH such as those at a cold CO₂ seep in Papua New Guinea exhibited lowered species diversity, structural complexity and resilience (Fabricius *et al.*, 2011).

Elevated sea temperature and increased precipitation are two impacts that have occurred in the recent past, the former at a global scale and the latter at a local scale. They give a fair warning of what can be expected and both impacts are examined further. The 1998 El Niño event provided a good opportunity to observe the effects of global warming. Elevated sea surface temperature occurred from mid-1997 to late 1998 and coral reefs displayed the most dramatic effect. Mass bleaching of corals took place worldwide at an unprecedented scale and highlighted the urgency of protecting reef resiliency. Management is thus needed to prevent compromise of reef system integrity. Mortality of shallow water corals was as high as 95% in some parts of the world while no mortality was observed in other places (Wilkinson & Hodgson, 1999).

Moderate to extensive bleaching was reported throughout Southeast Asia and all countries noted the extent of this event, which was unprecedented (Chou *et al.*, 2002). In Indonesia, bleaching started in early 1998 in West Sumatra resulting in over 90% mortality. It then spread to other reefs throughout the country causing decreases in live coral cover ranging from 30 to 90%. Recovery was variable after a few years with some reefs retaining depressed live coral cover of less than 10%, while for others it reached 40%. In the Philippines, mean live coral cover decreased by 19% after the 1998 bleaching in Tubbataha with no further loss or recovery after two years. At Danjungan Island in Negros Occidental where coral mortality from the bleaching was high, recovery was observed over the next two years. The species *Pavona clavus* recovered better in medium depths of 12 m compared to shallow waters of 6 m.

Widespread bleaching of shallow reefs in the Gulf of Thailand affected also the coral recruits. Corals on pinnacles in deeper water (10-15 m) escaped the bleaching. Local extinctions of some *Acropora* species were recorded while *Goniopora* showed complete recovery. Recovery in the inner Gulf of Thailand took a longer time because of low coral recruitment, but the east and west coasts of the Gulf had large numbers of coral recruits that facilitated recovery. In Vietnam's Con Dao islands, 37% of coral colonies bleached. Recovery was reported to be slow over the next two years. In Singapore, widespread mass bleaching occurred as sea surface temperature remained unusually high from January 1998, reaching 34.3°C in June. All hard coral species bleached, together with some species of soft corals and colonial sea anemones. However, sea surface temperature returned to normal (29.5 to 31.5°C) after June, allowing the bleached corals to recover and limit mortality to 20%.

Wilkinson and Hodgson (1999) noted that the 1997/98 bleaching event was the most severe ever observed and raised the question of whether this was just an isolated event or that similar events will follow at greater frequency as global warming continues. In the early part of 2010, the seas warmed up to temperatures higher than in 1997/98 and triggered widespread bleaching once again. Investigations revealed that species that were severely impacted in the 1997/98 bleaching appeared to be less impacted this time, while those that showed little effect to the earlier event were heavily bleached. The differences in species response and mortality patterns suggested some adaptive ability to thermal stress by coral species (Guest *et al.*, 2012).

Increased frequency of extreme weather is expected from global warming. Periods of drought interspersed by intense precipitation will cause wide salinity fluctuations in shallow waters and affect intertidal life. The impact of sudden lowering of salinity on intertidal biological communities was observed at a location in Singapore after unusually heavy rainfall over many decades occurred at the end of 2006 and early 2007 (Chou, 2010).

Excessive discharge from Malaysia's Johor River northeast of Singapore caused a sustained decline of salinity in that part of the Johor Strait. Chek Jawa, a protected intertidal habitat on the island of Pulau Ubin was fully exposed to the freshwater outflow for weeks. Mass mortality of sessile and sedentary osmotic-conforming species was observed and many species such as carpet anemones, sea stars and sea cucumbers, unable to regulate the osmotic concentration of their body fluids against the lowered salinity, literally burst open as the less saline water infused their body cavities. The intertidal flat was scattered with the fragmented remains of these species.

Investigations showed that recovery was good within a year after the event. Many species that suffered the mass kill had reappeared in abundance. At the same time, some species, which were not common before, became more abundant. These findings indicated that biodiversity can generally recover as environmental conditions revert to normal but long-term changes in the community structure are inevitable as indicated by the new appearance of the invasive Asian mussel, *Musculista senhousia*, a native of northern Southeast Asia that has spread to Singapore, Australia, New Zealand, the United States and the Mediterranean. In Singapore, *Musculista senhousia* established only in degraded habitats and its opportunistic colonization of Chek Jawa after the excessive rainfall in December 2007 demonstrated the potential of an invasive species to dominate a biodiversity-rich natural habitat following an abrupt environmental change and disrupt full recovery to the original community structure.

MANAGEMENT OPTIONS

The region's economic growth over the last four decades has had an impact on the coastal and marine habitats that are the richest in the world. Degraded habitats cannot provide the

high levels of ecosystem services that adequately protected habitats can and there are interesting challenges to sustainable development of the region's coastal and marine environment. Studies have indicated that ecosystems that are effectively managed can generate revenues from the ecosystem services of tourism and recreation. The annual income derived from Indonesia's Pulau Weh Marine Protected Area is US\$230,000, Malaysia's Pulau Redang \$545,000, Vietnam's Hon Mun MPA \$243,190 and Philippine's Lingayan Gulf \$4.7 million (CI, 2008).

In a cost-benefit analysis of coral mining for lime production in Lombok, coral miners could enjoy profits of US\$27,000/km² of reef but the societal cost in terms of fisheries and tourism loss from the destroyed reefs ranged from US\$33,000 to 762,000 (Öhman & Cesar, 2000). Hodgson and Dixon (1988) estimated that logging in Palawan could yield gross revenues of \$8.6 m over a 10-year span but at a cost to the fisheries (\$6.2 m) and tourism (\$13.9 m) industries dependent on the environment that will be impacted by the logging activity. Based on this analysis, the Philippine national government banned logging in Palawan and declared El Nido bay a marine reserve.

One of the most well-documented, long-term investigations of the impact of marine reserves on reef fish density was conducted at Sumilon and Apo Islands (Russ & Alcala, 2003). They demonstrated that fish yield was sustained from eleven years of continuous protection at Apo Island, while at Sumilon Island, the intermittent removal of protection resulted in significant declines of fish yield much greater than the build-up of fish biomass and density over the years under protection.

DISCUSSION

Climate change together with elevated sea surface temperature and ocean acidification are looming threats to the region's marine biodiversity. It will test the resilience of marine ecosystems, which are already compromised by intense anthropogenic pressure. The capacity of marine ecosystems to adapt to climate change impacts is severely undermined by present human demands that lead to excessive overexploitation, degradation, fragmentation and pollution. These pressures erode ecological integrity and depress ecosystem services.

Ecosystem resilience needs to be strengthened to minimise further damage from climate change impacts (SCBD, 2010) and the immediate focus is on the reduction of anthropogenic pressure and more efficient management of marine biodiversity. This is one of the great challenges facing the region, which has relied heavily on the services of marine ecosystems but not doing enough to maintain sustainability.

A sensible investment is to improve efforts at increasing the resiliency of ecosystems. This can be done through better management that includes more effective protection and restoration of degraded habitats such as seagrass, mangroves and coral reefs. There is much scope for research into reef restoration (Edwards & Gomez, 2007) and many reef restoration techniques have been initiated (Chou *et al.*, 2009). They include low cost approaches to hasten reef recovery from blast fishing where rubble stabilization and rock piles were found to encourage better coral recruitment and growth compared to scattered rubble. (Fox *et al.*, 2005; Raymundo *et al.*, 2007).

Against the gloomy outlook of habitat destruction are successful protection and rehabilitation measures at local levels (Tun *et al.*, 2008). Demonstration sites on reversing degradation trends have been established under different regional projects such as the UNEP/GEF South China Sea project (UNEP, 2004). Seeking to improve management for better sustainability, demonstration sites on more efficient integrated management of the

coast that addresses a sustainable development strategy for the seas of East Asia are being scaled up (PEMSEA, 2003).

REFERENCES

- ASEAN. 2006. *Third ASEAN State of the Marine Environment Report 2006*. ASEAN Secretariat, Jakarta.
- Barber, P. H., Erdmann M. V. & Palumbi S. R. 2006. Comparative phylogeography of three codistributed stomatopods: origins and timing of regional lineage diversification in the coral triangle. *Evolution*, **60(9)**: 1825-1839.
- Bellwood, D. R. & Meyer, C. P. 2009. Searching for heat in a marine biodiversity hotspot. *Journal of Biogeography*, **36**: 569-576.
- Burke, L., Reytar K., Spalding M. & Perry A. 2011. *Reefs at Risk Revisited*. World Resources Institute, Washington D.C.
- Chou, L. M. 1992. Potential Impacts of Climate Change and Sea-level Rise on the East Asian Seas Region. *Journal of Southeast Asian Earth Sciences*, **7(1)**: 61-64.
- Chou, L. M. (ed) 1994. *Implications of Expected Climate Changes in the East Asian Seas Region: An Overview*. RCU/EAS Technical Report Series No. 2. Regional Coordinating Unit, East Asian Seas, UNEP (Bangkok).
- Chou, L. M. 1996. Importance of Southeast Asian Marine Ecosystems in Biodiversity Maintenance. In Biodiversity and the Dynamics of Ecosystems (Turner, I. M., Diong C. H., Lim S. S. L. & Ng P. K. L. eds.), *DIWPA Series Volume. 1*: 227-235.
- Chou, L. M. 2010. *Anticipated Impacts of Climate Change on Marine Biodiversity: Using Field Situations that Simulate Climate Change in Singapore*. In *Moving Forward, Southeast Asian Perspectives on Climate Change and Biodiversity* (Sajise P. E., Ticsay M. V. & Saguiguit Jr. G. C. eds.), Southeast Asian Regional Center for Graduate Study and Research in Agriculture and Institute of Southeast Asian Studies, Singapore: 131-138.
- Chou, L. M., Vo, S. T., PhilReefs, Yeemin, T., Cabanban, A., Suharsono & Kessna, I. 2002. *Status of Southeast Asia Coral Reefs*. In *Status of Coral Reefs of the World: 2002* (Wilkinson C. ed.), Australian Institute of Marine Science, Townsville: 123-152.
- Chou, L. M., Yeemin, T., Abdul Rahim, G. Y., Vo, S. T., Alino, P. & Suharsono, S. 2009. Coral Reef Restoration in the South China Sea. *Galaxea*, **11**: 67-74.
- CI (Conservation International). 2008. *Economic Values of Coral reefs, Mangroves and Seagrasses: a Global Compilation*. Center for Applied Biodiversity Science, Conservation International.
- Djoghla, A. & Ganapin Jr, D. J. 2010. *Exploring the link between climate change and biodiversity*. In *Moving Forward: Southeast Asian Perspectives on Climate Change and Biodiversity* (Sajise P. E., Ticsay M. V., Saguiguit Jr G. C. eds.), Institute of Southeast Asian Studies and Southeast Asian Regional Center for Graduate Study and Research in Agriculture, Singapore: 3-8.
- Edwards, A. & Gomez, E. 2007. *Reef Restoration Concepts and Guidelines: Making Sensible Management Choices in the Face of Uncertainty. Coral Reef Targeted Research & Capacity Building for Management Programme*. St Lucia, Australia.
- FAO. 2007. *The World's Mangroves 1980-2005*. Food and Agriculture Organisation, Rome.
- Fabricius, K. E., Langdon, C., Uthicke S., Humphrey C., Noonan S., De'ath G, Okazaki R., Muehlllehner N., Glas M. S. & Janice M. L. 2011. Losers and Winners in Coral Reefs Acclimatized to Elevated Carbon Dioxide Concentrations. *Nature Climate Change*, **1**: 165-169.

- Fox, H. E., Mous, P. J., Pet, J. S., Muljadi, A. H. & Caldwell, R. L. 2005. Experimental Assessment of Coral Reef Rehabilitation Following Blast Fishing. *Conservation Biology*, **19**(1): 98-107.
- Green, E. P. & Short, F. T. (eds.) 2003. *World Atlas of Seagrasses*. World Conservation Monitoring Centre. University of California Press.
- Guest, J. R., Baird, A. H., Maynard, J. A., Muttaqin, E., Edwards, A. J., Cambell, S. J., Yewdall, K., Affendi Y.A. & Chou L. M. 2012. Contrasting Patterns of Coral Bleaching Susceptibility in 2010 Suggest an Adaptive Response to Thermal Stress. *PLoS One*, **7**(3): 705-713.
- Hodgson, G. & Dixon L. A. 1988. *Logging Versus Fisheries and Tourism in Palawan*. Occasional Papers No 7. East-West Environment and Policy Institute, Honolulu.
- Hughes, T. P., Baird, A. H., Bellwood, D. R., Card, M., Connolly, S. R., Folke, C., Grosberg, R., Hoegh-Guldberg, O., Jackson, J. B. C., Kleypas, J., Lough, J. M., Marshall, P., Nystrom, M., Palumbi, S. R., Pandolfi, J. M., Rosen, B. & Roughgarden, J. 2003. Climate Change, Human Impacts, and the Resilience of Coral Reefs. *Science*, **301**: 929-933.
- IPCC. 2007. *Climate Change 2007: Synthesis Report. Contribution of Working Groups I, II and III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change*. IPCC, Geneva, Switzerland.
- IUCN/UNEP. 1985. *Management and Conservation of Renewable Marine Resources in the East Asian Seas Region*. UNEP Regional Seas Reports and Studies No. 65. United Nations Environment Programme.
- JEC (ed.). 2000. *The State of the Environment in Asia 1999/2000*. Japan Environmental Council. Springer-Verlag Tokyo.
- Kelleher, G., Bleakley, C. & Wells, C. 1995. *A Global Representative System of Marine Protected Areas*. Volume 3. The World Bank, Washington D.C.
- Knittweis, L., Kraemer, W. E., Timm, J. & Kochzius, M. 2009. Genetic Structure of *Heliofungia actiniformis* (Scleractinia: Fungiidae) Populations in the Indo-Malay Archipelago: Implications for Live Coral Trade Management Efforts. *Conservation Genetics*, **10**: 241-249.
- Kochzius, M. & Nuryanto, A. 2008. Strong Genetic Population Structure in the Boring Giant Clam, *Tridacna crocea*, Across the Indo-Malay Archipelago: Implications Related to Evolutionary Processes and Connectivity. *Molecular Ecology*, **17**: 3775-3787.
- Kochzius, M., Seidel, C., Hauschild, J., Kirchhoff, S., Mester, P., Meyer-Wachsmuth, I., Nuryanto, A. & Timm, J. 2009. Genetic Population Structures of the Blue Starfish *Linckia laevigata* and its Gastropod Ectoparasite *Thyca crystallina*. *Marine Ecology Progress Series*, **396**: 211-219.
- Lourie, S. A. & Vincent, A. C. J. 2004. A Marine Fish Follows Wallace's Line: the Phylogeography of the Three-spot Seahorse (*Hippocampus trimaculatus*, Syngnathidae, Teleostei) in Southeast Asia. *Journal of Biogeography*, **31**(12): 1975-1985.
- MEA (Millennium Ecosystem Assessment) 2005. *Ecosystem and Human Well-being: Current Status and Trends Assessment*. Chapter 19 – Coastal systems: 515-549.
- Öhman, M. C. & Cesar, H. S. J. 2000. *Costs and Benefits of Coral Mining*. In *Collected Essays on Economics of Coral Reefs*, (Cesar H. S. J. Ed.). CORDIO, Department for Biology and Environmental Sciences, Kalmar University, Kalmar, Sweden: 85 - 93.
- PEMSEA (Partnerships in Environmental Management for the Seas of East Asia) 2003. *Sustainable Development Strategy for the Seas of East Asia: Regional Implementation of the World Summit on Sustainable Development Requirements for the Coasts and Oceans*. PEMSEA, Quezon City, Philippines.

- Rao, A. N. 1986. *Mangrove Ecosystems of Asia and the Pacific*. In: *Mangroves of Asia and the Pacific: Status and Management. Technical Report of the UNDP/UNESCO Research and Training Pilot Programme on Mangrove Ecosystems in Asia and the Pacific (RAS/79/002)*. Natural Resources Management Centre and National Mangrove Committee, Ministry of Natural Resources, Philippines: 1-48.
- Raymundo, L. J., Maypa, A. P., Gomez, E. D. & Cadiz, P. 2007. Can Dynamite-blasted Reefs Recover? A Novel, Low-tech Approach to Stimulating Natural Recovery in Fish and Coral Populations. *Marine Pollution Bulletin*, **54(7)**: 1009-1019.
- Roberts, C. M., McClean, C. J., Veron, J. E. N., Hawkins, J. P., Allen, G. R., McAllister, D. E., Mittermeier, C. G., Schueler, F. W., Spalding, M., Wells, F., Vynne, C., Werner, T. B. 2002. *Marine biodiversity hotspots and conservation priorities for tropical reefs*. *Science*, **295**: 1280-1284.
- Rogeli, R., Hare, B., Nabel, J., Macey, K., Schaeffer, M., Markmann, K. & Meinschausen M. 2009. Halfway to Copenhagen, No Way to 2⁰C. *Nature Reports Climate Change*, **3**: 81-83.
- Russ, G. R. & Alcala, A. C. 2003. Marine Reserves: Rates and Patterns of Recovery and Decline of Predatory Fish, 1983-2000. *Ecological Applications*, **13(6)**: 1553–1565.
- SCBD (Secretariat of the Convention on Biological Diversity). 2010. *Global Biodiversity Outlook 3*. Montreal.
- Spalding M., Ravilious C., Green E. P. 2001. *World atlas of coral reefs*. World Conservation Monitoring Centre. University of California Press.
- Silverman, J., Lazar, B., Cao, L., Caldeira, K. & Erez, J. 2009. Coral Reefs May Start Dissolving When Atmospheric CO₂ Doubles. *Geophysical Research Letters*. **36**: L05606.
- Singh, H. R., Chong, V .C., Sasekumar, A. & Lim, K. H. 1994. *Value of Mangroves as Nursery and Feeding Grounds*. In Proceedings, Third ASEAN-Australia Symposium on Living Coastal Resources, 16-20 May 1994, Bangkok (Wilkinson, C. R., Sudara, S. & Chou, L. M. eds.), Australian Institute of Marine Science, Townsville: 105-116.
- Timm, J. & Kochzius, M. 2008. Geological History and Oceanography of the Indo-Malay Archipelago Shape the Genetic Population Structure in the False Clown Anemonefish (*Amphiprion ocellaris*). *Molecular Ecology*, **17(18)**: 3999-4014.
- Tun, K., Chou, L. M., Thamasak, Y., Phongsuwan, N., Affendi, Y. A., Ho, N., Sour, K., Nguyen, V. L., Nanola, C., Lane, D. & Tuti, Y. 2008. *Status of Coral Reefs in Southeast Asia*. In *Status of Coral Reefs of the World: 2008* (Wilkinson, C. R. ed.), Australian Institute of Marine Science, Townsville: 131-144.
- UNEP. 2001. *Asia-Pacific Environmental Outlook*. United Nations Environment Programme, Bangkok.
- UNEP. 2004. *Coral Reefs in the South China Sea*. UNEP/GEF/SCS Technical Publication No. 2. UNEP/GEF Project Co-ordinating Unit, United Nations Environment Programme, Bangkok.
- UNEP. 2007. *Procedure for Determination of National and Regional Economic Values for Ecotone Goods and Services, and Total Economic Values of Coastal Habitats in the Context of the UNEP/GEF Project Entitled: "Reversing Environmental Degradation Trends in the South China Sea and Gulf of Thailand"*. South China Sea Knowledge Document No. 3. UNEP/GEF/SCS/Inf. 3.
- UNEP/COBSEA. 2010. *State of the Marine Environment Report for the East Asian Seas 2009* (Chou, L. M. ed.). COBSEA Secretariat, Bangkok.
- Weeks, R., Russ, G. R., Alcala, A. C. & White, A. T. 2010. Effectiveness of Marine Protected Areas in the Philippines for Biodiversity Conservation. *Conservation Biology*, **24(2)**: 531-540.

- Wilkinson, C. & Hodgson, G. 1999. Coral Reefs and the 1997-1998 Mass Bleaching and Mortality. *Nature & Resources*, **35(2)**: 16-25.
- Yap, H. T. 1994. *Implications of Expected Climatic Changes on Natural Coastal Ecosystems in the East Asian Seas Region*. In *Implications of Expected Climate Change in the East Asian Seas Region: An Overview* (Chou, L. M. ed), RCU/EAS Technical Report Series No. 2. Regional Coordinating Unit, East Asian Seas, UNEP (Bangkok): 105-122.