

HANDBOOK ON HABITAT RESTORATION

GENERAL PRINCIPLES AND CASE STUDIES IN SINGAPORE

EDITORS LENA CHAN / DANIEL NG / LIM LIANG JIM



© 2023 National Parks Board

Published by National Parks Board

Singapore Botanic Gardens, 1 Cluny Road, Singapore 259569

Website: www.nparks.gov.sg

All rights reserved.

No part of this publication may be reproduced, stored in a retrieval system, or transmitted in any form or by any means, electronic, mechanical, photocopying, recording or otherwise, without the prior permission of the copyright owners.

All images unless otherwise credited belong to the National Parks Board. All photographs may not be used without the written permission of the copyright holders.

ISBN 978-981-18-8480-1

Citation: Chan L, Ng D & Lim LJ (2023) Handbook on Habitat Restoration: General Principles and Case Studies in Singapore. National Parks Board, Singapore. 340 pp.

To learn more about our City in Nature, scan the QR code below or visit nparks.gov.sg/CityInNature



Share your love for nature and animals at
#NParksBuzz #AnimalBuzzSG

Cover design by: Nur Anisah Binte Mahmud

Typesetting by: Colin Lee & Amanda Yap

Front cover: Hampstead Wetlands Park (top); Eco-Link@BKE (bottom left); Rifle Range Nature Park (bottom right)

Back cover: Multi-tiered planting of native trees and shrubs along roadside (top left); Bishan-Ang Mo Kio Park (top right); Old Mandai Road (center); Red-Wattled Lapwings at Kranji Marshes (bottom left); Jurong Lake Gardens (bottom right)

CHAPTER 26

Reef Restoration in Singapore

Chou Loke Ming

The loss of more than half of Singapore's coral reef habitats to land reclamation and exposure of the remaining reefs to chronic sedimentation necessitate active restoration interventions. Singapore's reef restoration activities stretch back to the late 1980s. Two initiatives dealt with artificial reefs involving large and heavy structures that required barges and cranes to transport and deploy. The majority dealt with restoration techniques that could be handled by scuba-diving researchers. The earlier artificial reef project in 1989 deployed structures made from tyre pyramids and hollow concrete frames. The project was carried out by the National University of Singapore and supported under the ASEAN-USAID Coastal Resources Management Project. The scale of this artificial reef was dwarfed by the more recent artificial reef project (Singapore's largest artificial reef structures) in 2018 where 10-metre-tall purpose-built concrete and fibreglass structures were placed at Sisters' Islands Marine Park (See Chapter 14). This project was a collaboration between JTC Corporation (JTC) and National Parks Board (NParks).

The large majority of reef restoration projects dealt with coral translocation and strategies to enhance survival and growth of corals, and where structures were involved, they were small and portable. Collectively, the research has revealed much information valuable to the advancement of reef restoration in Singapore's marine environment. This would not have been possible without the funding support from and/or collaboration with various agencies, such as Keppel Group, Housing and Development Board, JTC, Maritime and Port Authority, National Environment Agency, NParks, National Research Foundation (Marine Science Research and Development Programme), Sentosa Development Corporation, Singapore Maritime Institute, Singapore Tourism Board, Wildlife Reserves Singapore Conservation Fund, and the European Union.

Reef restoration efforts employing techniques that overcome sedimentation challenges have indicated their viability (Ng *et al.*, 2016a; Chou *et al.*, 2018). These range from the provision of sloping solid substrata (Low *et al.*, 2006) or horizontal mesh surfaces that prevent sediment accumulation (Ng & Chou, 2014). This will complement the observed natural coral colonisation of seawalls (Chou *et al.*, 2010; Ng *et al.*, 2012) and other human-made structures such as jetty pilings (Chou & Lim, 1986; Ong & Tan, 2012) lining the urban coast.

In-situ coral nurseries have a significant role in reef restoration efforts in a sediment-stressed environment. Studies on two scleractinian species, *Pachyseris speciosa* and *Pocillopora acuta* (Poquita-Du *et al.*, 2017) indicated that fragments raised in nurseries for five months before being transplanted to the reef substrate, grew significantly faster by three to five times compared to those that were transplanted directly without a nursery phase (Afiq-Rosli *et al.*, 2017). The faster growth of transplants in nurseries augmented their size and continued to manifest after final transplantation to the reef, enabling them to perform better than directly transplanted fragments.

Nurseries can also be used to nurture ‘corals of opportunity’ (COPs), which are naturally fragmented corals lying free on the reef floor or coral juveniles that have recruited on loose rubble. COPs raised in nurseries at Pulau Semakau (Chou *et al.*, 2009) and nubbins (small fragments) raised in nurseries at St John’s and Lazarus Islands (Bongiorni *et al.*, 2011) showed that improved survivorship and growth rate can be achieved in Singapore’s sediment-impacted waters. These studies highlight the feasibility of establishing coral nurseries in other locations to preserve scleractinian genotypes from reefs that will be directly affected by coastal development.

The coral nurseries themselves served as microhabitats to enhance biodiversity despite the sedimentation, attracting a large variety of species such as fishes (Taira *et al.*, 2016) and reef-associated invertebrates (Wee *et al.*, 2019). These nurseries also provided opportunities for recruitment, settlement, and development of reef fauna. They enhance ecosystem functioning in degraded as well as non-reefal sites while nurturing coral fragments for transplantation (Chou *et al.*, 2018).

It is important to consider preserving the locality’s genetic diversity when transplanting corals from a source reef about to be exposed to development impact. Investigations of four coral species from such a reef showed that 33% to 40% of colonies can represent 80% of genetic diversity and more than 50% of colonies represent more than 90% of genetic diversity (Afiq-Rosli *et al.*, 2019). Hence, not all colonies need to be relocated when resources are constrained. Colonies of hermaphroditic species can be collected from over a smaller area, while gonochoric species can be collected over a larger spatial area, the latter having greater genetic variability over larger distances.

Biodiversity on artificial substrates developed through natural colonisation can be enhanced by active transplantation particularly in the intertidal zone. Natural colonisation showed the viability

of hard structures such as seawalls as habitats for biological communities (Ng *et al.*, 2015) and active transplantation can further enhance the structure's ecological value. Fragments of five species of scleractinian corals, three species of soft corals and three species of sponges reared in *ex-situ* nurseries prior to transplantation exhibited variable survivorship and growth. The scleractinian coral *Porites lobata*, soft coral *Lobophytum* sp., and sponge *Lendenfeldia chondrodes* had high survivorship, rapid growth and extended mean survival times two years after transplantation. Coral species with massive or encrusting growth forms were best at establishing and developing on seawalls and observed to provide shelter and food for reef fish and gastropods.

Active restoration can help to increase the intertidal biodiversity of seawalls but must take into account that not all species can survive conditions at seawalls, particularly those with an early developing pioneering but competitive community. Similarly, not all species do well when transplanted to subtidal seawalls. Of six species investigated, *Hydnophora rigida*, *Podabacia crustacea*, *Echinopora lamellosa* and *Platygyra sinensis* had sustained growth and survival rates exceeding 90% after six months (Toh *et al.*, 2017). The study also showed how involvement of volunteers could lower the labour cost of the transplantation effort. In a recent transplantation project from 2013 to 2019, 904 coral fragments were transplanted to cover 150 square metres of degraded reef and create 272 square metres of 'new' reefs (Chou *et al.*, 2016). The 'new' reefs are non-reefal areas on which coral transplantation was attempted to determine if reef communities could be generated.

Many restoration initiatives are supported by short-term projects over durations that do not allow for long-term assessment. Sessile lifeform diversity developed from natural recruitment on fibreglass reef enhancement units (Ng *et al.*, 2016b) increased significantly over a span of 10 years with hard corals and coralline algae contributing most to the temporal dissimilarity. The reef units augmented ecosystem functioning with 119 sessile and mobile taxa utilising them for food and shelter. It must be emphasised that long-term monitoring is essential for assessing the effectiveness of reef restoration efforts (Chou *et al.*, 2016).

References

- Afiq-Rosli L, Huang D, Toh TC, Taira D, Ng CSL, Song TC & Chou LM (2019) Maximising genetic diversity during coral transplantation from a highly impacted source reef. *Conservation Genetics*, 20: 629–637.

- Afiq-Rosli L, Taira D, Loke HX, Toh TC, Toh KB, Ng CSL, Cabaitan PC, Chou LM & Song TC (2017) In situ nurseries enhance coral transplant growth in sedimented waters. *Marine Biology Research*, 13(8): 878–887.
- Borngiorni L, Giovanelli D, Rinkevich B, Pusceddu A, Chou LM & Danovaro R (2011) First step in the restoration of a highly degraded coral reef (Singapore) by in situ coral intensive farming. *Aquaculture*, 322–323: 191–200.
- Chou LM & Lim TM (1986) A preliminary study of the coral community on artificial and natural substrates. *Malayan Nature Journal*, 39: 225–229.
- Chou LM, Ng CSL, Chan SMJ & Seow LA (2010) Natural colonization of a marina seawall in Singapore. *Journal of Coastal Development*, 14(1): 11–17.
- Chou LM, Sam SQ, Toh TC & Ng CSL (2018) The relevance of reef restoration in the continually changing marine environment of Singapore. *Vietnam Journal of Marine Science and Technology*, 18(3): 278–285.
- Chou LM, Toh TC & Ng CSL (2016) Effectiveness of reef restoration in Singapore's rapidly urbanizing coastal environment. *International Journal of Environmental Science and Development*, 8(8): 576–580.
- Chou LM, Yeemin T, Abdul Rahim BGY, Vo ST, Alino P & Suharsono (2009) Coral reef restoration in the South China Sea. *Galaxea, Journal of Coral Reef Studies*, 11(2): 67–74.
- Low TL, Tanzil JTI & Chou LM (2006) Preliminary study of community development and scleractinian recruitment on fiberglass artificial reef units in the sedimented waters of Singapore. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 16(1): 61–76.
- Ng CSL, Chen D & Chou LM (2012) Hard coral assemblages on seawalls in Singapore. In: Tan KS (ed.) *Contributions to Marine Science*, Tropical Marine Science Institute, National University of Singapore, pp. 75–79.
- Ng CSL & Chou LM (2014) Rearing juvenile 'corals of opportunity' in in-situ nurseries – a reef rehabilitation approach for sediment-impacted environments. *Marine Biology Research*, 10(8): 833–838.
- Ng CSL, Lim SC, Ong JY, Teo LMS, Chou LM, Chua KE & Tan KS (2015) Enhancing the biodiversity of coastal defence structures: transplantation of nursery-reared reef biota onto intertidal seawalls. *Ecological Engineering*, 82: 480–486.
- Ng CSL, Toh TC & Chou LM (2016a) Coral restoration in Singapore's sediment-challenged sea. *Regional Studies in Marine Science*, 8(3): 422–429.

- Ng CSL, Toh TC & Chou LM (2016b) Artificial reefs as a reef restoration strategy in sediment-affected environments: Insights from long-term monitoring. *Aquatic Conservation: Marine and Freshwater Ecosystems*, 27(5): 976–985.
- Ong JIJ & Tan KS (2012) Observations on the subtidal fouling community on jetty pilings in the southern islands of Singapore. In: Tan KS (ed.) *Contributions to Marine Science*, Tropical Marine Science Institute, National University of Singapore, pp. 121–126.
- Poquita-Du R, Ng C, Loo J, Afiq-Rosli L, Tay Y, Todd P, Chou LM & Huang D (2017) New evidence shows that *Pocillopora* ‘damicornis-like’ corals in Singapore are actually *Pocillopora acuta* (Scleractinia: Pocilloporidae). *Biodiversity Data Journal*, 5: e11407.
- Taira D, Toh TC, Ng CSL, Sam SQ & Chou LM (2016) Coral nurseries as habitats for juvenile corallivorous butterflyfish. *Marine Biodiversity*, 47(3): 787–788.
- Toh TC, Ng CSL, Loke HX, Taira D, Toh KB, Afiq-Rosli L, Poquita-Du R, Cabaitan P, Sam SQ, Kikuzawa YP, Chou LM & Song TC (2017) A cost-effective approach to enhance scleractinian diversity on artificial shorelines. *Ecological Engineering*, 99: 349–357.
- Wee SYC, Sam SQ, Sim WT, Ng CSL, Taira D, Afiq-Rosli L, Kikuzawa YP, Toh TC & Chou LM (2019) The role of in situ coral nurseries in supporting mobile invertebrate epifauna. *Journal for Nature Conservation*, 50: 125710.