

ENVIRONMENT-INDUCED COLOUR CHANGES IN THE MASSIVE CORALS *DIPLOASTREA SPECIOSA* AND *DIPLOASTREA HELIOPORA*

Peter A. Todd^{1*}, Wanting Zhao¹ and Loke Ming Chou²

¹Experimental Marine Ecology Laboratory, Department of Biological Sciences
National University of Singapore, 14 Science Drive 4, Singapore 117557, Republic of Singapore

²Reef Ecology Laboratory, Department of Biological Sciences
National University of Singapore, 14 Science Drive 4, Singapore 117557, Republic of Singapore

(*Corresponding author: dbspat@nus.edu.sg)

ABSTRACT. — Environment-dependent variation in the morphological, physiological or behavioural expression of a genotype is termed phenotypic plasticity. Twelve colonies of both *Diploastrea speciosa* and *Diploastrea heliopora*, two common Indo-Pacific massive corals, were tested for plastic responses by transplanting palm-sized fragments (clonemates) over six stations (three reefs × two depths) in Singapore's southern waters. After 15 weeks, there were no major changes in colour among the fragments transplanted to the shallow stations, with the typical grey of *Diploastrea heliopora*, and the bright green oral discs of the *Diploastrea speciosa* polyps, clearly represented. The fragments at the two deepest stations, however, changed dramatically. The oral discs of *Diploastrea speciosa* lost their bright green pigmentation and the surrounding tissue turned pale brown. The response from *Diploastrea heliopora* was even more pronounced, with yellow-pinks and pale greens replacing its usual grey pigmentation. Plasticity in colour was evident for both species, and probably light-induced, however, further work is required to determine whether such changes are adaptive.

KEY WORDS. — acclimatisation, colour, depth, phenotypic plasticity, reef, sediment, Singapore

INTRODUCTION

Even though it is well known that corals exhibit a wide range of colours within and among species, their function remains largely unexplored (Takabayashi & Hoegh-Guldberg, 1995; Salih et al., 1998). The most vivid colours, the brilliant greens, blues, and reds, are host (animal) based and related to Green Fluorescent Proteins (GFPs), probably pocilloporins (Dove et al., 2001; Shagin et al., 2004; Oswald et al., 2007; Gruber et al., 2008). These are generally found in the epithelial cells of the coral polyp (Salih et al., 2000; Mazel et al., 2003), usually above the zooxanthellae in high light environments and below them in low light environments (Schlichter et al., 1994; Salih et al., 1998; Dove et al., 2001). They have also been found in close association with zooxanthellae in the gastroderm (Oswald et al., 2007). In addition, a family of non-fluorescent proteins known as chromoproteins (CPs) contributes to the bright purple and blue colours displayed by some species (Dove et al., 2001; Shagin et al., 2004). The brown colour common to most corals is attributed to their algal symbiont: zooxanthellae (Hochberg et al., 2003). The algal pigments enable the zooxanthellae to utilise light of different wavelengths for photosynthesis. Of the spectrum that makes up photosynthetically active radiation (PAR), 400–550 nm is absorbed by chlorophyll *a*, chlorophyll *c*, and the carotenoid peridinin, whereas 650–700 nm is absorbed almost entirely by chlorophyll *a* (Falkowski et al., 1990). The host pigments are believed to have at least two, non-exclusive, purposes, i.e., to protect the coral from harmful ultra-violet radiation (UVR) (Salih et al., 2000; Dove et al., 2001), and to absorb UVR and re-emit the energy at longer wavelengths that can be photosynthesised (Schlichter & Fricke, 1990). It has also been proposed that GFPs have a role in attracting zooxanthallae, which are drawn to green light (Hollingsworth et al., 2005).

Short UVR wavelengths are damaging to coral reef epifauna (Jokiel, 1980). Of the ultra-violet spectrum, UV-A (320–400 nm) is considered the least deleterious, UV-B (280–320 nm) is highly biocidal, but the majority of the lethal UV-C (200–280 nm) does not reach Earth's surface (Jokiel & York, 1982). Corals taken from deep water, where less UVR penetrates, tend to be more sensitive to UVR than those acclimatised to the shallows (Falkowski et al., 1990). Elevated UVR causes greater photosynthetic activity and production of O₂, which results in a proportionate increase in production of reactive oxygen species (ROS) (Jamieson et al., 1986). These ROS collectively lead to a condition known as photo-oxidative stress, causing damage to DNA, membrane proteins, enzymes, lipids, and photosystem components (Lesser, 2006), eventually causing coral bleaching (Downs et al., 2002). UVR photo-inhibition is a result of chloroplast damage (Jokiel & York, 1982) and photo-pigment destruction (Wood, 1987). GFPs are thought to be part of a broad antioxidant defence system as they neutralise O₂, a ROS (Bou-abdallah et al., 2006). Further, GFPs may also directly convert excess UVR into radiation at wavelengths less detrimental to the zooxanthellae (Dove et al., 2001). They are, however, not physical barriers to UVR (D'Angelo et al., 2008).

As the variation in both host and zooxanthellae pigment types and quantities can often be correlated to light character and intensity, they are believed to be photo-adaptive (Jokiel & York, 1982; Chalker et al., 1983; Titlyanov, 1991; Salih et al., 1998). Light is directly linked to the expression of pigment proteins, causing either an increase or decrease in concentration of different pigments when intensity is increased (D'Angelo et al., 2008). Jokiel & York (1982) found more near-UV absorbing substances (identified as mycosporinlike amino acids [MAAs] by Dunlap & Chalker [1979]) in *Pocillopora damicornis* in high UVR environments. The green and brown morphs of *Porites astreoides* (Caribbean coral) have different depth distributions with the green morphs, more abundant in shallow (≤ 2 m) waters, better able to endure UVR (Gleason, 1993). The green morphs have more MAAs which block 310–360 nm but are transparent to PAR. Total concentrations of MAAs decreased with depth in 10 species of coral from Hawaii and Belize (Banaszak et al., 1998), and UVR had a significant positive effect on the quantity of MAAs present in colonies of the coral *Porites compressa* (Kuffner, 2001).

It has also been suggested that other animal pigments absorb UVR and fluoresce the energy as longer wavelengths suitable for photosynthesis (Falkowski et al., 1990), possibly giving the host coral a competitive edge in low light environments (Schlichter et al., 1994). Wavelength transference in the deep-water (95–145 m) coral *Leptoseris fragilis* is believed to help this species survive in extremely low light conditions (Schlichter & Fricke, 1990; Kaiser et al., 1993). In 17 out of 71 zooxanthellate species, Schlichter et al. (1994) detected autofluorescent chromatophores transforming 365–410 nm wavelengths into 430–500 nm. Zooxanthellae and the autofluorescent chromatophores are often found densely packed in the oral area (Schlichter & Fricke, 1990; Schlichter et al., 1994). However, Gilmore et al. (2003) and Mazel et al. (2003) disagree that GFPs enhance photosynthesis, as there is no observable photon transfer between GFPs and zooxanthellae. Although widespread, the specific function that GFP serves in the coral host has yet to be determined conclusively (Mazel et al., 2003; Field et al., 2006; Oswald et al., 2007).

In zooxanthellae, pigmentation usually increases in each cell with depth, but the actual number of cells can increase or decrease (Falkowski et al., 1990; Kaiser et al., 1993; Iglesias-Prieto & Trench, 1994). Zooxanthellae densities decreased with depth while content of pigment per cell increased in *Leptoseris fragilis* (Kaiser et al., 1993), *Fungia repanda*, and *Fungia echinata* (Masuda et al., 1993). In the pocilloporid *Stylophora pistillata*, the density of zooxanthellae and their pigmentation concentration both increased under 30% and 8% surface PAR, but at 0.8% surface PAR zooxanthellae were lost through degradation (Titlyanov et al., 2001). Zooxanthellae with higher levels of pigment become more efficient light harvesters at greater depth, but why their density should change is less clear (Titlyanov et al., 2001).

Even though Salih et al. (1998) and Schlichter et al. (1994) explain where pigments are concentrated in the coral tissue, few authors describe in detail how study corals actually look in differing environments, or before and after transplantation. During a large study to test for morphological plasticity in two massive coral species: *Diploastrea speciosa* and *Diploastrea heliophora* (Todd et al., 2004), dramatic changes in pigmentation were observed. Photographs of these changes were published by Todd et al. (2002a), but with very little accompanying explanatory text. Here we discuss in greater depth, the colour changes, their probable causes, and their potential functional significance.

MATERIAL AND METHODS

The methods for the transplant part of this study are described in full in Todd et al. (2004). In brief: in May 2000, fragments of *Diploastrea speciosa* (formerly *Favia speciosa*; Budd et al., 2012) and *Diploastrea heliophora* were reciprocally transplanted among three reefs: Cyrene Reef, Pulau Hantu, and Raffles Lighthouse. Cyrene Reef is a submerged reef close to Singapore Island (4 km), heavily impacted by sediment originating from nearby dredging of shipping lanes, reclamation activities, and river discharge. Pulau Hantu is a small island, 7.5 km from Singapore Island. As a result of the many surrounding islands and reefs, Pulau Hantu has the calmest waters of the three study reefs. Raffles Lighthouse is situated 15 km offshore with open sea to the south and is the least affected by sedimentation (but possibly exposed to more wave action).

Two stations were established at each reef, one deep and one shallow (the depths of each station are provided in Fig. 1). Four colonies of both species were sampled randomly from each of the three reefs. From every colony, six fragments (clone-mates) were excised and distributed randomly throughout the six stations (3 reefs \times 2 depths). At each station, 24 fragments (12 of each species), with their tissue-covered surface facing upwards, were attached to rectangular aluminium racks at a height of ~ 0.45 m above the substrate. Fifteen weeks after the corals fragments were moved to their new habitats, 2:1 photographs were taken of all specimens in situ and with a Nikonos V camera loaded with Fujichrome 'Velvia' colour slide film (ASA 50). Flash lighting and distance from the coral was fixed for all photographs taken. The resultant images were grouped by genotype and examined for phenotypic change.




Fig. 1. Fragments (clone-mates) of *Diploastrea speciosa* and *Diploastrea heliophora* were reciprocally transplanted across six stations (3 reefs \times 2 depths). The depth (below mean sea level) of each station is provided on the y-axis. These representative images were taken 15 weeks later. For each pair of images, *Diploastrea speciosa* is always on the left and *Diploastrea heliophora* is always on the right. Note: For each species, all six fragments originated from the same donor colony.

RESULTS

After 15 weeks of transplantation, the shallow water fragments at Cyrene were most similar to how *Diploastrea speciosa* and *Diploastrea heliophora* appeared in their 'natural' environment. There was no obvious change in pigmentation, with the typical grey of *Diploastrea heliophora*, and the bright green oral discs of the *Diploastrea speciosa* polyps, clearly represented (Fig. 1). The fragments at the deep station, however, changed dramatically. The oral discs of *Diploastrea speciosa* lost their bright green pigmentation and the surrounding tissues turned pale brown. The response from *Diploastrea heliophora* was even more pronounced, with very little of its characteristic grey pigmentation remaining. The coral also appeared to have lost its undulating topography.

At Pulau Hantu (shallow station), the lateral and vertical expansion of polyps discerned by the morphometric analysis of cleaned skeletons by Todd et al. (2004) was clearly visible (Fig. 1). Furthermore, six buds could be seen growing in between mature *Diploastrea heliophora* polyps, as described by Todd et al. (2002b). The topography of this fragment also appeared rugged when compared to Cyrene Reef (shallow station). At the deep station, the oral discs of *Diploastrea speciosa* were only distinguishable from the surrounding tissue by their shape in relief, not by their colour. The polyps appeared lacklustre. Similar to Cyrene Reef (deep station), the tissue of *Diploastrea heliophora* had a translucent quality.

For the fragments transplanted to the shallow station at Raffles Lighthouse, the polyps of both species, and particularly those of *Diploastrea heliophora*, were more exert than at Cyrene Reef (shallow station) and pigmentation was strong for both species (Fig. 1). At the 7 m station (the least deep and least sediment-impacted of the deep stations), the loss of pigmentation was not so dramatic compared to the other deep sites, for example, the green oral discs of *Diploastrea speciosa* were still visible and colours in the *Diploastrea heliophora* fragments were more pronounced.

DISCUSSION

The timing and nature of this survey enabled observations to be made that were not feasible later on as, by the end of 30 weeks when the experiment was terminated, many of the deep fragments at Cyrene Reef and Pulau Hantu were severely degraded, making comparisons of all six cogeners impossible. The images presented here, taken after 15 weeks and representative of the overall results, illustrate both physiological and morphological plasticity in *Diploastrea speciosa* and *Diploastrea heliophora* (Todd et al., 2002a), with changes in colour especially clear. The changes in morphology that occurred have been described in detail elsewhere (Todd et al., 2004; Todd, 2008), therefore, only the induced changes in pigmentation will be considered here.

Kawaguti (1937) suspended clone-mates of *Pavona praetora* at 1-m intervals to a depth of 10 m in the sea near Palao, South Pacific (underwater visibility 6–7 m). After three weeks the specimens had turned yellowish near the surface, reaching a maximum in brown colouration at 4 m and becoming paler again between 4–10 m. The pattern of colouration observed by Kawaguti (1937) is similar to that of *Diploastrea speciosa* in the present study if the deep station at Raffles Lighthouse is considered an intermediate site for light (being shallower and in less sedimented waters

than the other two deep sites). Fig. 1 illustrates how the fragments transplanted to 7 m at Raffles Lighthouse were darker than those moved to shallow and deep stations elsewhere. As an increase in UVR and PAR reduces the photosynthetic capabilities of symbiotic algae (Lesser, 2000), the amount of photosynthetic pigments and therefore intensity of these browns usually decreases with increased UVR and PAR (Oswald et al., 2007; Torres et al., 2007). Correspondingly, fragments of *Diploastrea speciosa* at shallow stations appeared less brown compared to the fragments from the deep stations.

Lewis (1974) found juveniles of *Favia fragum* were almost colourless after 12 weeks of dark incubation and Miller (1995) noted that whereas exposed colonies of *Oculina arbuscula* were brown, those living under overhangs were white. These findings are analogous to the changes in *Diploastrea heliopora*, which displayed an inverse relationship between pigmentation and depth (Fig. 1), i.e., a mid-grey colouration in the shallows, changing to light ruddy-brown at Raffles Lighthouse (deep station), finally turning to pale yellow-pink, with inter-polyp areas of mid-green, at the greatest depths. Titlyanov et al. (2001) found, as a result of zooxanthellae degradation, a net loss of chlorophyll per polyp for *Stylophora pistillata* in extremely low light (0.8%). A similar loss of zooxanthellae in *Diploastrea heliopora* and *Diploastrea speciosa* may explain the paleness of fragments transplanted to the deep stations at Cyrene Reef and Pulau Hantu, where conservative estimates of light penetration to are, respectively, 1.13% and 0.58% surface PAR (Todd et al., 2004). Being located slightly shallower and with greater light penetration, the fragments at Raffles Lighthouse (deep station) displayed a darker brown colour compared to colonies at Cyrene and Pulau Hantu, which had distinctive pale green inter-polyp areas (Fig. 1). The increasing intensity of these green lines with decreasing light seems to suggest a photo-enhancement role to assist zooxanthellae photosynthesis under reduced light conditions. The cause for the loss of zooxanthellae with increasing depth is less clear, but may be because the zooxanthellae type hosted by the coral could not survive at such low light levels.

At Raffles Lighthouse (deep station), *Diploastrea speciosa* appeared to be gaining pigmentation whereas *Diploastrea heliopora* seemed to be losing it. This may have been related to their different depths of origin. Around Singapore, *Diploastrea speciosa* is common between 3–6 m, but *Diploastrea heliopora* is more frequently found in depths of <3 m. Therefore, the deep station at Raffles Lighthouse, with irradiance levels of only ~1.55% surface PAR (Todd et al., 2004), may be more detrimental to *Diploastrea heliopora* than *Diploastrea speciosa*. Contrary to this argument, at the termination of this experiment (15 weeks after these images were taken) *Diploastrea heliopora* appeared to have experienced less tissue loss than *Diploastrea speciosa* at all the deep stations.

Diploastrea speciosa around Singapore is characterised by its bright green oral disc (probably due to GFPs). If the pigment causing this colouration aided photosynthesis through auto-fluorescence, then it could be expected to be more apparent, or at least equally apparent, in low light. Instead, the intensity of the green diminished with depth, suggesting that it is an UVR absorbing and/or reflecting pigment that is redundant in deeper waters where UVR levels were reduced. Only the deep station at Raffles Lighthouse still supported *Diploastrea speciosa* with green oral discs, again probably due to more light penetrating here than the deep stations at Cyrene Reef and Pulau Hantu.

Since the sites differed primarily in the amount of light reaching the fragments, it would seem that pigment levels in *Diploastrea speciosa* and *Diploastrea heliopora* are regulated by light intensity. For both species, there was a change in colouration from predominantly beige (for *Diploastrea speciosa*) and grey (for *Diploastrea heliopora*), to various intensities of brown, yellow and green. The increase in depth may have reduced the expression of certain host pigments responsible for the colours at the shallow stations, possibly because of reduced need for protection from UVR, thereby allowing the brown colour of the symbionts to be displayed more prominently. Zooxanthellae density may also have affected the intensity of the colours exhibited. It is not known, however, whether the concentration of photo-pigments in zooxanthellae increased, or only became visible due to the absence of host pigments masking their colours. It would be insightful to know the quantities of each pigment, and the density of zooxanthellae, in the various clone-mates before and after transplantation but, unfortunately, these data were not collected. Plasticity in colour is evident in both species and probably light-induced, however, further work is required to determine whether such changes are adaptive.

ACKNOWLEDGEMENTS

Many thanks to Peta Sanderson, Roy Sidle, and James Guest for their editing, discussion, and help in the field. This work was funded by grant numbers RP 3991014 and R-154-000-414-133.

LITERATURE CITED

- Banaszak, A. T., M. P. Lesser, I. B. Kuffner & M. Ondrusek, 1998. Relationship between ultraviolet (UV) radiation and mycosporinlike amino acids (MAAs) in marine organisms, *Bulletin of Marine Science*, **63**: 617–628.
- Bou-Abdallah, F., N. D. Chasteen & M. P. Lesser, 2006. Quenching of superoxide radicals by green fluorescent protein.

- Biochimica et Biophysica Acta*, **1760**: 1690–1695.
- Budd, A. F., H. Fukami, N. D. Smith & N. Knowlton, 2012. Taxonomic classification of the reef coral family Mussidae (Cnidaria: Anthozoa: Scleractinia). *Zoological Journal of the Linnean Society*, **166**: 465–529.
- Chalker, B. E., W. E. Dunlap & J. K. Oliver, 1983. Bathymetric adaptations of reef-building corals at Davies Reef, Great Barrier Reef, Australia. II. Light saturation curves for photosynthesis and respiration. *Journal of Experimental Marine Biology and Ecology*, **73**: 37–56.
- D'Angelo, C., A. Denzel, A. Vogt, M. V. Matz, F. Oswald, A. Salih, G. U. Nienhaus & J. Wiedenmann, 2008. Blue light regulation of host pigment in reef-building corals. *Marine Ecology Progress Series*, **364**: 97–106.
- Dove, S. G., O. Hoegh-Guldberg & S. Ranganathan, 2001. Major colour patterns of reef-building corals are due to family of GFP-like proteins. *Coral Reefs*, **19**: 197–204.
- Downs, C. A., J. E. Fauth, J. C. Halas, P. Dustan, J. Bemiss & C. M. Woodley, 2002. Oxidative stress and seasonal coral bleaching. *Free Radical Biology and Medicine*, **33**: 533–543.
- Dunlap, W. C. & B. E. Chalker, 1986. Identification and quantification of near-UV absorbing compounds (S-320) in a hermatypic scleractinian. *Coral Reefs*, **5**: 155–159.
- Falkowski, P., P. L. Jokiel & R. A. Kinzie, 1990. Irradiance and corals. In: Dubinski, Z. (ed.), *Ecosystems of the World 25: Coral Reefs*. Elsevier, Amsterdam. Pp. 89–107.
- Field, S. F., M. Y. Bulina, I. V. Kelmanson, J. P. Bielawski & M. V. Matz, 2006. Adaptive evolution of multicolored fluorescent proteins in reef-building corals. *Journal of Molecular Evolution*, **62**: 332–339.
- Gilmore, A. M., A. W. D. Larkum, A. Salih, S. Itoh, Y. Shibata, C. Bena, H. Yamasaki, M. Papina & R. Van Woesik, 2003. Simultaneous time resolution of the emission spectra of fluorescent proteins and zooxanthellar chlorophyll in reef-building corals. *Photochemistry and Photobiology*, **77**: 515–523.
- Gleason, D. F., 1993. Differential effects of ultraviolet radiation on green and brown morphs of the Caribbean coral *Porites astreoides*. *Limnology and Oceanography*, **38**: 1452–1463.
- Gruber, D. F., H. Kao, S. Janoschka, J. Tsai & V. A. Pieribone, 2008. Patterns of fluorescent protein expression in scleractinian corals. *Biology Bulletin*, **215**: 143–154.
- Hochberg, E. J., M. J. Atkinson & S. Andréfouët, 2003. Spectral reflectance of coral reef bottom-types worldwide and implications for coral reef remote sensing. *Remote Sensing of Environment*, **85**: 159–173.
- Hollingsworth, L. L., R. A. Kinzie, T. D. Lewis, D. A. Krupp & J. A. C. Leong, 2005. Phototaxis of motile zooxanthellae to green light may facilitate symbiont capture by coral larvae. *Coral Reefs*, **24**: 523–523.
- Iglesias-Prieto, R. & R. K. Trench, 1994. Acclimation and adaptation to irradiance in symbiotic dinoflagellates. I. Responses of the photosynthetic unit to changes in photon flux density. *Marine Ecology Progress Series*, **113**: 163–175.
- Jamieson, D., B. Chance, E. Cadenas & A. Boveris, 1986. The relation of free radical production to hyperoxia. *Annual Review of Physiology*, **48**: 703–719.
- Jokiel, P. L., 1980. Solar ultra-violet radiation and coral reef epifauna. *Science*, **207**: 1069–1071.
- Jokiel, P. L. & R. H. Jr. York, 1982. Solar ultraviolet photobiology of the reef coral *Pocillopora damicornis* and symbiotic zooxanthellae. *Bulletin of Marine Science*, **32**: 301–315.
- Kaiser, P., D. Schlichter & H. W. Fricke, 1993. Influence of light on algal symbionts and the deep water coral *Leptoseris fragilis*. *Marine Biology*, **117**: 45–52.
- Kawaguti, S., 1937. On the physiology of reef corals III. Regeneration and phototropism in reef corals. *Palao Tropical Biological Station Studies*, **2**: 209–218.
- Kuffner, I. B., 2001. Effects of ultraviolet (UV) radiation on larval settlement of the reef coral *Pocillopora damicornis*. *Marine Ecology Progress Series*, **217**: 251–261.
- Lesser, M. P., 2000. Depth-dependent photoacclimatization to solar ultraviolet radiation in the Caribbean coral *Montastraea faveolata*. *Marine Ecology Progress Series*, **192**: 137–151.
- Lesser, M. P., 2006. Oxidative stress in marine environments: Biochemistry and physiological ecology. *Annual Review of Physiology*, **68**: 253–278.
- Lewis, J. B., 1974. The settlement behavior of planulae larvae of the hermatypic coral *Favia fragum* (Esper). *Journal of Experimental Biology and Ecology*, **15**: 165–172.
- Masuda, K., M. Goto, T. Maruyama & S. Miyachi, 1993. Adaptation of solitary corals and their zooxanthellae to low light and UV radiation. *Marine Biology*, **117**: 685–691.
- Mazel, C. H., M. P. Lesser, M. Y. Gorbunov, T. M. Barry, J. H. Farrell, K. D. Wyman & P. G. Falkowski, 2003. Green-fluorescent proteins in Caribbean corals. *Limnology and Oceanography*, **48**: 402–411.
- Miller, M. W., 1995. Growth of a temperate coral: Effects of temperature, light, depth, and heterotrophy. *Marine Ecology Progress Series*, **122**: 217–225.
- Oswald, F., F. Schmitt, A. Leutenegger, S. Ivanchenko & C. D'Angelo, 2007. Contributions of host and symbiont pigments to the coloration of reef corals. *Federation of European Biochemical Societies Journal*, **274**: 1102–1122.
- Salih, A., O. Hoegh-Guldberg & G. Cox, 1998. Photoprotection of symbiotic dinoflagellates by fluorescent pigments in reef corals. In: Greenwood, J. G. & N. J. Hall (eds.), *Proceedings of the Australian Coral Reef Society 75th Anniversary Conference*. Pp. 217–230.
- Salih, A., A. Larkum, G. Cox, M. Kühl & O. Hoegh-Guldberg, 2000. Fluorescent pigments in corals are photoprotective. *Nature*, **408**: 850–853.

- Shagin, D. A., E. V. Barsova, Y. G. Yanushevich, A. F. Fradkov, K. A. Lukyanov, Y. A. Labas, T. N. Semenova, J. A. Ugalde, A. Meyers, J. M. Nunez, E. A. Widder, S. A. Lukyanov & M. V. Matz, 2004. GFP-like proteins as ubiquitous metazoan superfamily: Evolution of functional features and structural complexity. *Molecular Biology Evolution*, **21**: 841–850.
- Schlichter, D., U. Meier & H. W. Fricke, 1994. Improvement of photosynthesis in zooxanthellate corals by autofluorescent chromatophores. *Oecologia*, **99**: 124–131.
- Schlichter, D. & H. W. Fricke, 1990. Coral host improves photosynthesis of endosymbiotic algae. *Naturwissenschaften*, **77**: 447–450.
- Takabayashi, M. & O. Hoegh-Guldberg, 1995. Ecological and physiological differences between two colour morphs of the coral *Pocillopora damicornis*. *Marine Biology*, **123**: 705–714.
- Titlyanov, E. A., 1991. The stable level of coral primary production in a wide light range. *Hydrobiologia*, **216**: 383–387.
- Titlyanov, E. A., T. V. Titlyanova, K. Yamazato & R. van Woesik, 2001. Photo-acclimation dynamics of the coral *Stylophora pistillata* to low and extremely low light. *Journal of Experimental Marine Biology and Ecology*, **263**: 211–225.
- Todd, P. A., R. C. Sidle & L. M. Chou, 2002a. Plastic corals from Singapore 2. *Coral Reefs*, **21**: 407–408.
- Todd, P. A., R. C. Sidle & L. M. Chou, 2002b. Plastic corals from Singapore 1. *Coral Reefs*, **21**: 391–392.
- Todd, P. A., R. J. Ladle, N. J. I. Lewin-Koh & L. M. Chou, 2004. Genotype × environment interactions in transplanted clones of the massive corals *Favia speciosa* and *Diploastrea heliopora*. *Marine Ecology Progress Series*, **271**: 167–182.
- Todd, P. A., 2008. Morphological plasticity in scleractinian corals. *Biological Reviews*, **83**: 315–337.
- Torres, J. L., R. A. Armstrong, J. E. Corredor & F. Gilbes, 2007. Physiological responses of *Acropora cervicornis* to increased solar irradiance. *Photochemistry and Photobiology*, **83**: 839–850.
- Wood, W. F., 1987. Effect of solar ultra-violet radiation on the kelp *Ecklonia radiata*. *Marine Biology*, **96**: 143–150.